

On Hamiltonian minimal submanifolds in the space of oriented geodesics in real space forms

Nikos Georgiou*, G. A. Lobos

29 November 2014

Abstract

We prove that a deformation of a hypersurface in a $(n+1)$ -dimensional real space form $\mathbb{S}_{p,1}^{n+1}$ induce a Hamiltonian variation of the normal congruence in the space $\mathbb{L}(\mathbb{S}_{p,1}^{n+1})$ of oriented geodesics. As an application, we show that every Hamiltonian minimal submanifold in $\mathbb{L}(\mathbb{S}^{n+1})$ (resp. $\mathbb{L}(\mathbb{H}^{n+1})$) with respect to the (para-) Kähler Einstein structure is locally the normal congruence of a hypersurface Σ in \mathbb{S}^{n+1} (resp. \mathbb{H}^{n+1}) that is a critical point of the functional $\mathcal{W}(\Sigma) = \int_{\Sigma} (\prod_{i=1}^n |\epsilon + k_i^2|)^{1/2}$, where k_i denote the principal curvatures of Σ and $\epsilon \in \{-1, 1\}$. In addition, for $n = 2$, we prove that every Hamiltonian minimal surface in $\mathbb{L}(\mathbb{S}^3)$ (resp. $\mathbb{L}(\mathbb{H}^3)$) with respect to the (para-) Kähler conformally flat structure is locally the normal congruence of a surface in \mathbb{S}^3 (resp. \mathbb{H}^3) that is a critical point of the functional $\mathcal{W}'(\Sigma) = \int_{\Sigma} \sqrt{H^2 - K + 1}$ (resp. $\mathcal{W}'(\Sigma) = \int_{\Sigma} \sqrt{H^2 - K - 1}$), where H and K denote, respectively, the mean and Gaussian curvature of Σ .

1 Introduction

The space $\mathbb{L}(M)$ of oriented geodesics of a pseudo-Riemannian manifold (M, g) has been of great interest for the last three decades and has been studied by different authors (see for example [1],[2],[5],[6],[7],[8],[14],[15]). When (M, g) is a Riemannian symmetric space of rank one, Alekseevsky, Guilfoyle and Kligenberg have described in [1] all possible metrics defined on $\mathbb{L}(M)$ that are invariant under the isometry group of g .

*The author is partially supported by Fapesp (2010/08669-9).

In the case where (M, g) is a real $(n + 1)$ -dimensional space form $\mathbb{S}_{p,1}^{n+1}$ of signature $(p, n+1-p)$ with constant sectional curvature one, Anciaux has showed in [2] that $\mathbb{L}(\mathbb{S}_{p,1}^{n+1})$ admits a Kähler or a para-Kähler structure (G, \mathbb{J}, Ω) , where \mathbb{J} is the complex or paracomplex structure and Ω is the symplectic structure such that the metric G is Einstein and is invariant under the isometry group of g . In the same work, for $n = 2$, Anciaux has proved that $\mathbb{L}(\mathbb{M})$ admits an extra Kähler or para-Kähler structure $(G', \mathbb{J}', \Omega)$, where \mathbb{J}' is the complex or paracomplex structure, such that the invariant metric G' is of neutral signature, locally conformally flat and is invariant under the isometry group of g .

The submanifold theory of $\mathbb{L}(M)$ gives some interesting informations about the submanifold theory of M . For example, the normal congruence (or Gauss map) of a one-parameter family of parallel hypersurfaces in M is a Lagrangian submanifold (the induced symplectic structure identically) of the corresponding space of geodesics (see for example [2]). In particular, the normal congruence $L(\Sigma)$ of a Weingarten surface Σ in $\mathbb{S}_{p,1}^3$ (its principal curvatures are functionally related) is flat with respect to the metric G' induced on $L(\Sigma)$ [2].

Let (M, J, g, ω) be a (para-) Kähler manifold and let $\phi : \Sigma \rightarrow M$ be a Lagrangian immersion. A normal vector field X is called *Hamiltonian* if $X = J\nabla u$, where J is the (para-) complex structure and ∇u is the gradient of $u \in C^\infty(\Sigma)$ with respect to the non-degenerate induced metric ϕ^*g . We say that a variation (ϕ_t) of ϕ is a *Hamiltonian variation* if its velocity $X = \partial_t|_{t=0}\phi_t$ is a Hamiltonian vector field with the additional condition that the function u is compactly supported. The Lagrangian immersion ϕ is said to be *Hamiltonian minimal* or *H-minimal* if it is a critical point of the volume functional with respect to Hamiltonian variations. The first variation formula of the volume functional implies that a Hamiltonian minimal submanifold is characterised by the equation $\text{div} JH = 0$, where H denotes the mean curvature vector of ϕ and div is the divergence operator with respect to the induced metric [11]. Further study of *H-minimal* submanifolds can be found at the following articles [4, 9, 10, 12].

Palmer showed in [13] that a smooth variation of a hypersurface in the sphere \mathbb{S}^{n+1} induces a Hamiltonian variation of the Gauss map in $\mathbb{L}(\mathbb{S}^{n+1})$. An analogue result for the 3-dimensional Euclidean space \mathbb{E}^3 has been shown by Anciaux, Guilfoyle and Romon in [3]. Following similar computations that Palmer used in [13], we prove that any smooth variation of a hypersurface in the real space form $\mathbb{S}_{p,1}^{n+1}$ induces a Hamiltonian variation in the symplectic manifold $(L^\pm(\mathbb{S}_{p,1}^{n+1}), \Omega)$. In particular we prove the following:

THEOREM 1.1. *Let $\phi_t, t \in (-\epsilon, \epsilon)$ be a smooth one-parameter deformation of*

an immersion $\phi := \phi_{t=0}$, of the n -dimensional oriented manifold Σ in the real space form $\mathbb{S}_{p,1}^{n+1}$. Then, the corresponding Gauss maps Φ_t form a Hamiltonian variaton with respect to the symplectic manifold $(L^\pm(\mathbb{S}_{p,1}^{n+1}), \Omega)$.

Anciaux, Guilfoyle and Romon have proved in [3] that a Hamiltonian minimal surface in $\mathbb{L}(\mathbb{E}^3)$ (resp. $\mathbb{L}(\mathbb{E}_1^3)$) is the Gauss map of a surface S in \mathbb{E}^3 (resp. \mathbb{E}_1^3) which is a critical point of the functional $\mathcal{F} = \int_S \sqrt{H^2 - K} dA$, where H and K denote the mean curvature and the Gauss curvature, respectively. In this article we extend this result for the case of the space $\mathbb{L}(\mathbb{S}_{p,1}^{n+1})$ of oriented geodesics in a $(n+1)$ -dimensional real space form. In particular, we consider the (para-) Kähler Einstein structure (G, \mathbb{J}, Ω) and the locally conformally flat (para-) Kähler structure $(G', \mathbb{J}', \Omega)$ both endowed on $\mathbb{L}(\mathbb{S}_{p,1}^{n+1})$. Then, as an application of Theorem 1.1, we prove the following:

THEOREM 1.2. *Let $\phi : \Sigma^n \rightarrow \mathbb{S}_{p,1}^{n+1}$ be a real diagonalizable hypersurface in $\mathbb{S}_{p,1}^{n+1}$ and let Φ be the Gauss map of ϕ . Then, away of umbilic points, we have the following statements:*

(i) *The Gauss map Φ is a Hamiltonian minimal submanifold with respect to the (para-)Kähler Einstein structure (G, \mathbb{J}) if and only if the immersion ϕ is a critical point of the functional*

$$\mathcal{W}(\phi) = \int_{\Sigma} \sqrt{\prod_{i=1}^n |\epsilon + k_i^2|} dV,$$

where k_1, \dots, k_n are the principal curvatures of ϕ and ϵ denotes the length of the normal vector field of ϕ .

(ii) *For $n = 2$, the Gauss map Φ is a Hamiltonian minimal surface in $(L^\pm(\mathbb{S}_{p,1}^3), G', \mathbb{J}')$ if and only if the surface ϕ is a critical point of the functional*

$$\mathcal{W}'(\phi) = \int_{\Sigma} |k_1 - k_2| dA,$$

where k_1, k_2 denote the principal curvatures of ϕ .

Acknowledgements. The authors would like to thank H. Anciaux and Martin A. Magid for their helpful and valuable suggestions and comments.

2 Preliminaries

For $n \geq 1$, consider the Euclidean space \mathbb{R}^{n+2} endowed with the canonical pseudo-Riemannian metric of signature $(p, n+2-p)$, where $0 \leq p \leq n+2$:

$$\langle \cdot, \cdot \rangle_p = - \sum_{i=1}^p dx_i^2 + \sum_{i=p+1}^{n+2} dx_i^2.$$

Define the $(n+1)$ -dimensional real space form

$$\mathbb{S}_{p,1}^{n+1} = \{x \in \mathbb{R}^{n+2} \mid \langle x, x \rangle_p = 1\},$$

and let $\iota : \mathbb{S}_{p,1}^{n+1} \hookrightarrow (\mathbb{R}^{n+2}, \langle \cdot, \cdot \rangle_p)$ be the canonical inclusion. The induced metric $\iota^* \langle \cdot, \cdot \rangle_p$ has signature $(p, n+1-p)$ and is of constant sectional curvature $K = 1$.

Following the notations of [2] we denote by $L^+(\mathbb{S}_{p,1}^{n+1})$ (resp. $L^-(\mathbb{S}_{p,1}^{n+1})$) the set of spacelike (resp. timelike) oriented geodesics of $\mathbb{S}_{p,1}^{n+1}$, that is,

$$L^\pm(\mathbb{S}_{p,1}^{n+1}) = \{x \wedge y \in \Lambda^2(\mathbb{R}^{n+2}) \mid y \in T_x \mathbb{S}_{p,1}^{n+1}, \langle y, y \rangle_p = \epsilon\},$$

where $\epsilon = 1$ (resp. $\epsilon = -1$) corresponds to $L^+(\mathbb{S}_{p,1}^{n+1})$ (resp. $L^-(\mathbb{S}_{p,1}^{n+1})$). If $\Lambda^2(\mathbb{R}^{n+2})$ is equipped with the flat pseudo-Riemannian metric:

$$\langle \langle x_1 \wedge y_1, x_2 \wedge y_2 \rangle \rangle = \langle x_1, x_2 \rangle_p \langle y_1, y_2 \rangle_p - \langle x_1, y_2 \rangle_p \langle x_2, y_1 \rangle_p,$$

we denote by G the metric $\langle \langle \cdot, \cdot \rangle \rangle$ induced by the inclusion map $i : L^\pm(\mathbb{S}_{p,1}^{n+1}) \hookrightarrow \Lambda^2(\mathbb{R}^{n+2})$. On the other hand, in $L^\pm(\mathbb{S}_{p,1}^{n+1})$ can be defined a complex (paracomplex) structure \mathbb{J} as follows:

Let J be the canonical complex (paracomplex) structure in the oriented plane $x \wedge y \in L^\pm(\mathbb{S}_{p,1}^{n+1})$ defined by $Jx = y$ and $Jy = -\epsilon x$. Thus, $J^2 = -\epsilon Id$. A tangent vector to $i(L^\pm(\mathbb{S}_{p,1}^{n+1}))$ at the point $x \wedge y$ is of the form $x \wedge X + y \wedge Y$, where $X, Y \in (x \wedge y)^\perp$ in $\Lambda^2(\mathbb{R}^{n+2})$. The complex (paracomplex) structure \mathbb{J} is defined by:

$$\mathbb{J}(x \wedge X + y \wedge Y) := (Jx) \wedge X + (Jy) \wedge Y = y \wedge X + \epsilon x \wedge Y.$$

The metric G and the (para) complex structure \mathbb{J} are invariant under the natural action of the isometry group of $\mathbb{S}_{p,1}^{n+1}$. For $n \geq 3$, it has been showed in [1] that G is the unique invariant metric under the natural action of $SO(n+2-p, p)$.

The 2-form Ω defined by $\Omega(\cdot, \cdot) = \epsilon G(\mathbb{J}\cdot, \cdot)$, is a symplectic structure on $L^\pm(\mathbb{S}_{p,1}^{n+1})$ and in particular:

PROPOSITION 2.1. [2] *The quadruple $(L^+(\mathbb{S}_{p,1}^{n+1}), G, \mathbb{J}, \Omega)$ is a $2n$ -dimensional Kähler manifold with signature $(2p, 2n-2p)$, while $(L^-(\mathbb{S}_{p,1}^{n+1}), G, \mathbb{J}, \Omega)$ is a $2n$ -dimensional para-Kähler manifold. In both cases, the metric G is Einstein with constant scalar curvature $S = 2\epsilon n^2$.*

We now consider the case of $L^\pm(\mathbb{S}_{p,1}^3) \subset \Lambda^2(\mathbb{R}^4)$. The orthogonal $(x \wedge y)^\perp$ of an oriented plane $x \wedge y \in L^\pm(\mathbb{S}_{p,1}^3)$, is also a plane in \mathbb{R}^4 and is oriented in such a way its orientation is compatible with the orientation of the plane $x \wedge y$. Then it is possible to define a canonical complex or paracomplex structure J' , depending of whether the metric $\langle\langle \cdot, \cdot \rangle\rangle$ induced on $(x \wedge y)^\perp$ is positive or indefinite. In this case, we may define a complex or paracomplex structure \mathbb{J}' on $L^\pm(\mathbb{S}_{p,1}^3)$ by

$$\mathbb{J}'(x \wedge X + y \wedge Y) := x \wedge (J'X) + y \wedge (J'Y).$$

The pseudo-Riemannian metric G' on $L^\pm(\mathbb{S}_{p,1}^3)$ is given by:

$$G'(\cdot, \cdot) := \Omega(\cdot, J'\cdot) = -\epsilon G(\cdot, \mathbb{J} \circ \mathbb{J}'\cdot).$$

Furthermore,

PROPOSITION 2.2. [2] *The quadruples $(L^\pm(\mathbb{S}_{p,1}^3), G', \mathbb{J}', \Omega)$ are 4-dimensional (para-) Kähler manifolds. The metric G' is of neutral signature $(2, 2)$, scalar flat, locally conformally flat and is invariant under the natural action of $SO(4-p, p)$.*

Let $\phi : \Sigma^n \rightarrow \mathbb{S}_{p,1}^{n+1}$ be an immersion of a n -dimensional orientable manifold into the real space form $\mathbb{S}_{p,1}^{n+1}$ and let N be the unit normal vector of the hypersurface $S = \phi(\Sigma)$. The set \bar{S} of geodesics that are orthogonal to S , oriented in the direction of N , is called *the normal congruence* or *the Gauss map* of S . Then,

PROPOSITION 2.3. [2] *Let ϕ be an immersion of an orientable manifold Σ^n in $\mathbb{S}_{p,1}^{n+1}$ with unit normal vector field N . Then the Gauss map of $S = \phi(\Sigma)$ is the image of the map $\Phi : \Sigma^n \rightarrow L^\pm(\mathbb{S}_{p,1}^{n+1})$ defined by $\Phi = \phi \wedge N$. When Φ is an immersion, it is Lagrangian. Conversely, let $\Phi : \Sigma^n \rightarrow L^\pm(\mathbb{S}_{p,1}^{n+1})$ be an immersion of a simply connected n -manifold. Then $\bar{S} := \Phi(\Sigma)$ is the Gauss map of an immersed hypersurface of $\mathbb{S}_{p,1}^{n+1}$ if and only if Φ is Lagrangian.*

3 Hamiltonian minimal submanifolds

Throughout the article, when we talk about hypersurfaces in $\mathbb{S}_{p,1}^{n+1}$ or about Lagrangian submanifolds in $L^\pm(\mathbb{S}_{p,1}^{n+1})$, we mean that the induced metric is non-degenerate.

3.1 Hamiltonian variations in $L^\pm(\mathbb{S}_{p,1}^{n+1})$

Consider the $2n$ -dimensional symplectic manifold $(L^\pm(\mathbb{S}_{p,1}^{n+1}), \Omega)$, where $L^\pm(\mathbb{S}_{p,1}^{n+1})$ denotes the space of oriented geodesics in the real space form $\mathbb{S}_{p,1}^{n+1}$. Then we prove our first main result:

PROOF OF THEOREM 1.1: Let $\phi_t : \Sigma^n \rightarrow \mathbb{S}_{p,1}^{n+1}$, where $t \in (-t_0, t_0)$ for some $t_0 > 0$, be a smooth variation of an immersion $\phi := \phi_0 : \Sigma^n \rightarrow \mathbb{S}_{p,1}^{n+1}$ of an oriented n -dimensional manifold Σ in $\mathbb{S}_{p,1}^{n+1}$. Let $S_t := \phi_t(\Sigma)$ be the hypersurfaces of $\mathbb{S}_{p,1}^{n+1}$ and $S := \phi(\Sigma)$.

We denote by N_t the 1-parameter family of vector fields such that $N_0 = N$ and $\langle N_t, \phi_t \rangle_p = 0$. Following the notation of [13], there exist a smooth function f on S and a smooth section Y of the tangent bundle TS such that

$$\dot{\phi} = fN + Y,$$

where $\dot{\phi} = \partial_t \phi_t|_{t=0}$. Derivating the expression $\langle N_t, \phi_t \rangle_p = 0$ with respect to t at $t = 0$, we have

$$\langle \dot{N}, \phi \rangle_p = -\langle N, \dot{\phi} \rangle_p.$$

Then, it yields

$$\langle \dot{N}, \phi \rangle_p = -\epsilon f,$$

where $\epsilon = \langle N, N \rangle_p$. For $X_t \in TS_t$, where $X_0 := X \in TS$, we also derive with respect to t the expression $\langle N_t, X_t \rangle_p = 0$ at $t = 0$, and we obtain

$$\langle \dot{N}, X \rangle_p = -\epsilon df(X) + \langle dN(Y), X \rangle_p,$$

which finally gives

$$\dot{N} = -\epsilon \nabla f + dN(Y) - \epsilon f \phi.$$

Let $\Phi : \Sigma^n \rightarrow L^\pm(\mathbb{S}_{p,1}^{n+1}) : x \mapsto \phi(x) \wedge N(x)$ be the Gauss map of the immersion $\phi : \Sigma^n \rightarrow \mathbb{S}_{p,1}^{n+1}$. If $\bar{X} := d\Phi(X)$, from [2], we have that

$$\bar{X} = X \wedge N + AX \wedge \phi.$$

Let $\dot{\Phi} = \partial_t \Phi_t|_{t=0}$ be the velocity of the variation Φ_t and write $\dot{\Phi} = \dot{\Phi}^\top + \dot{\Phi}^\perp$, where $\dot{\Phi}^\top$ and $\dot{\Phi}^\perp$ denote the tangential and the normal component of $\dot{\Phi}$ respectively. Then,

$$\begin{aligned} G(\dot{\Phi}, \mathbb{J}\bar{X}) &= G(Y \wedge N + \epsilon \nabla f \wedge \phi - dN(Y) \wedge \phi, \mathbb{J}(X \wedge N + AX \wedge \phi)) \\ &= -df(X), \end{aligned}$$

which implies that

$$\dot{\Phi}^\perp = -\mathbb{J}\bar{\nabla} f,$$

and this completes the Theorem. \square

3.2 Applications

A *real diagonalizable immersion* is a smooth immersion ϕ of a hypersurface Σ^n into the $(n+1)$ -dimensional real space form $\mathbb{S}_{p,1}^{n+1}$ such that, locally, the shape operator A can be diagonalized, that is, it is existed a local orthonormal frame (e_1, \dots, e_n) and real functions k_1, \dots, k_n where, $A = \text{diag}(k_1, \dots, k_n)$. In this case, each vector field e_i is called a *principal direction* with corresponded *principal curvature* k_i .

REMARK 3.1. Note that every hypersurface in the Riemannian case, away of umbilic points, is real diagonalizable.

We are now in position to prove our second result:

PROOF OF THEOREM 1.2: Consider a smooth immersion of ϕ of the n -dimensional manifold Σ in $\mathbb{S}_{p,1}^{n+1}$ and let $\Phi : \Sigma^n \rightarrow L^\pm(\mathbb{S}_{p,1}^{n+1})$ be the corresponded Gauss map. The fact that ϕ is real diagonalizable implies the existence of an orthonormal frame (e_1, \dots, e_n) , with respect to the induced metric ϕ^*g , such that

$$Ae_i = k_i e_i, \quad i = 1, \dots, n,$$

where A denotes the shape operator of ϕ . Let $(\phi_t)_{t \in (-t_0, t_0)}$ be a smooth variation of ϕ and (Φ_t) be the corresponded variation of the Gauss map Φ . Real diagonalizability implies that the minimal polynomial of A is the product of distinct linear factors. Using the fact that the variation (Φ_t) is at least C^1 -smooth, it is possible to obtain a positive real number $t_1 < t_0$ such that ϕ_t is real diagonalizable for every $t \in (-t_1, t_1)$. We may extend all extrinsic geometric quantities such as the shape operator A , the principal directions e_i and the principal curvatures k_i to the 1-parameter family of immersions (ϕ_t) .

(i) From [2], the induced metric Φ_t^*G is given by $\Phi_t^*G = \epsilon \phi_t^*g + \phi_t^*g(A, A)$ and thus,

$$\Phi_t^*G = \text{diag}\left(\epsilon_1(\epsilon + k_1^2), \dots, \epsilon_n(\epsilon + k_n^2)\right),$$

where $\epsilon_i = g(e_i, e_i)$. For every sufficiently small $t > 0$, the volume of every Gauss map Φ_t , with respect to the metric G , is

$$(1) \quad \text{Vol}(\Phi_t) = \int_{\Sigma} \sqrt{|\det \Phi_t^*G|} dV = \mathcal{W}(\phi_t).$$

If ϕ is a critical point of the functional \mathcal{W} , we have

$$\partial_t(\text{Vol}(\Phi_t)) = 0,$$

for any Hamiltonian variation of Φ . Therefore, Φ is a Hamiltonian minimal submanifold with respect to the Kähler Einstein structure (G, J) . The converse follows directly from (1).

(ii) Assume that $n = 2$. From [2], in terms of the orthonormal frame (e_1, e_2) , the induced metric Φ^*G' is

$$\Phi^*G' = \begin{pmatrix} 0 & \epsilon_2(k_2 - k_1) \\ \epsilon_2(k_2 - k_1) & 0 \end{pmatrix}.$$

where $\epsilon_2 = \phi^*g(e_2, e_2)$. Then, the volume of every Gauss map Φ_t , with respect to the metric G' , is

$$(2) \quad \text{Vol}'(\Phi_t) = \int_{\Sigma} \sqrt{|\det \Phi_t^*G'|} dV = \mathcal{W}'(\phi_t),$$

and thus the second statement of the Theorem follows by a similar argument with the proof of the first statement. \square

Let $\phi : \Sigma^n \rightarrow \mathbb{S}_{p,1}^{n+1}$ be a hypersurface and N denotes the unit normal vector field. For $\theta \in \mathbb{R}$, consider the immersion $\phi_\theta := \cos \epsilon(\theta)\phi + \sin \epsilon(\theta)N$, where $\epsilon := |N|^2$ and $(\cos \epsilon(\theta), \sin \epsilon(\theta)) = (\cos \theta, \cos \theta)$ if $\epsilon = 1$ while for $\epsilon = -1$ we have $(\cos \epsilon(\theta), \sin \epsilon(\theta)) = (\cosh \theta, \cosh \theta)$. The images $\phi_\theta(\Sigma)$ and $\phi(\Sigma)$ are called parallel hypersurfaces. It is important to mention that parallel hypersurfaces have the same Gauss map [2].

Looking more carefully the relations (1) and (2) we obtain the following symmetry for the functionals \mathcal{W} and \mathcal{W}' :

COROLLARY 3.2. *If ϕ_1 and ϕ_2 are parallel smooth real diagonalizable immersions of the n -manifold Σ in $\mathbb{S}_{p,1}^{n+1}$ then $\mathcal{W}(\phi_1) = \mathcal{W}(\phi_2)$. In the case where $n = 2$, we also have that $\mathcal{W}'(\phi_1) = \mathcal{W}'(\phi_2)$.*

Using the Einstein (para-) Kähler structure $(L^\pm(\mathbb{S}_{p,1}^3), G, J)$ we obtain the following Corollary:

COROLLARY 3.3. *Let $\phi : \Sigma^n \rightarrow \mathbb{S}_{p,1}^{n+1}$ be a real diagonalizable hypersurface and let k_1, \dots, k_n be the principal curvatures. If Φ is the Gauss map of ϕ , then the function $\sum_{i=1}^n \tan \epsilon^{-1}(k_i)$ is harmonic with respect to the induced metric Φ^*G if and only if ϕ is a critical point of the functional $\int_{\Sigma} \sqrt{\prod_{i=1}^n |\epsilon + k_i^2|}$.*

PROOF. Let Φ be the Gauss map of ϕ and consider the Einstein (para-) Kähler structure (G, J) . Then, from [2], we know that the mean curvature \vec{H} of Φ is given by

$$\vec{H} = \frac{\epsilon}{n} J \nabla \left(\sum_{i=1}^n \tan \epsilon^{-1}(k_i) \right),$$

where ∇ denotes the Levi-Civita connection of the induced metric Φ^*G . Then, the Corollary follows by the following relation,

$$\operatorname{div}(nJ\vec{H}) = \Delta\left(\sum_{i=1}^n \tan \epsilon^{-1}(k_i)\right),$$

where div and Δ denote the divergence operator and the Laplacian of Φ^*G . \square

Using the Remark 3.1, we obtain the following two Corollaries:

COROLLARY 3.4. *Let $\phi : \Sigma^n \rightarrow \mathbb{S}^{n+1}$ (resp. in the hyperbolic space \mathbb{H}^{n+1}) be a hypersurface in the sphere \mathbb{S}^{n+1} . Then the Gauss map Φ is a Hamiltonian minimal submanifold in the (para-) Kähler Einstein structure (G, \mathbb{J}) if and only if the hypersurface ϕ is a critical point of the functional $\mathcal{W}(\phi) = \int_{\Sigma} \sqrt{\prod_{i=1}^n (1 + k_i^2)}$ (resp. $\mathcal{W}(\phi) = \int_{\Sigma} \sqrt{\prod_{i=1}^n |1 - k_i^2|}$), where k_1, \dots, k_n are the principal curvatures of ϕ .*

COROLLARY 3.5. *Let $\phi : \Sigma \rightarrow \mathbb{S}^3$ (resp. in the hyperbolic space \mathbb{H}^3) be a surface in the sphere \mathbb{S}^3 . Then the Gauss map Φ is a Hamiltonian minimal submanifold with respect to the Kähler, conformally flat structure (G', \mathbb{J}') if and only if the surface ϕ is a critical point of the functional $\mathcal{W}'(\phi) = \int_{\Sigma} \sqrt{H^2 - K + 1}$ (resp. $\mathcal{W}'(\phi) = \int_{\Sigma} \sqrt{H^2 - K - 1}$), where H, K denote the mean and the Gauss curvature of ϕ .*

References

- [1] D. ALEKSEEVSKY, B. GUILFOYLE AND W. KLINGENBERG, *On the geometry of spaces of oriented geodesics*, Ann. Global Anal. Geom. **40** (2011), 389–409.
- [2] H. ANCIAUX, *Space of geodesics of pseudo-Riemannian space forms and normal congruences of hypersurfaces*, Trans. Amer. Math. Soc. **366** (2014), 2699–2718.
- [3] H. Anciaux, B. Guilfoyle, P. Romon, *Minimal submanifolds in the tangent bundle of a Riemannian surface*, J. Geometry and Physics. **61**, 237–247 (2011).
- [4] A. Butscher and J. Corvino, *Hamiltonian stationary tori in Kähler manifolds*, Calc. Var. Partial Differential Equations. **45**, 63–100 (2012).
- [5] N. Georgiou and B. Guilfoyle, *On the space of oriented geodesics of hyperbolic 3-space*, Rocky Mountain J. Math. **40**, 1183–1219 (2010).

- [6] B. Guilfoyle and W. Klingenberg, *An indefinite Kähler metric on the space of oriented lines*, J. London Math. Soc. **72**, 497–509 (2005).
- [7] B. Guilfoyle and W. Klingenberg, *A neutral kähler metric on the space of time-like lines in Lorentzian 3-space* (2005) math.DG/0608782.
- [8] N.J. Hitchin, *Monopoles and geodesics*, Comm. Math. Phys. **83**, 579–602 (1982).
- [9] D. Joyce, Y.-I. Lee and R. Schoen, *On the existence of Hamiltonian stationary Lagrangian submanifolds in symplectic manifolds*, Amer. J. Math. **133**, 1067–1092 (2011).
- [10] Y.-I. Lee, *The existence of Hamiltonian stationary Lagrangian tori in Kähler manifolds of any dimension*, Calc. Var. Partial Differential Equations **45**, 231–251 (2012).
- [11] Y.-G. Oh, *Second variation and stabilities of minimal lagrangian submanifolds in Kähler manifolds*, Invent. Math. **101**, 501–519 (1990)
- [12] Y.-G. Oh, *Volume minimization of Lagrangian submanifolds under Hamiltonian deformations*, Math. Z. **212**, 175–192 (1993).
- [13] B. Palmer, *Buckling eigenvalues, Gauss maps and Lagrangian submanifolds*, Diff. Geom. Appl., **4**, 391–403 (1994).
- [14] M. Salvai, *On the geometry of the space of oriented lines in Euclidean space*, Manuscripta Math. **118**, 181–189 (2005).
- [15] M. Salvai, *On the geometry of the space of oriented lines of hyperbolic space*, Glasg. Math. J. **49**, 357–366 (2007).

Nikos Georgiou, UFAM, Manaus, AM, Brazil.
 nikos@ime.usp.br

G. A. Lobos, UFSCar, São Carlos, SP, Brazil.
 lobos@dm.ufscar.br