

THE BAOUENDI-TREVES APPROXIMATION THEOREM FOR CONTINUOUS VECTOR FIELDS

S. BERHANU AND J. HOUNIE

ABSTRACT. This article establishes the Baouendi-Treves approximation theorem for locally integrable structures whose vector fields have continuous coefficients. As a consequence, some uniqueness results are derived.

1. INTRODUCTION

Let \mathcal{M} be a C^∞ manifold of dimension N and suppose \mathcal{L} is a continuous vector subbundle of $\mathbb{C}T(\mathcal{M})$ with fiber dimension $1 \leq n < N$ and set $m = N - n$. We will say that \mathcal{L} is locally integrable if every $p \in \mathcal{M}$ is contained in an open set Ω such that there exist m functions $Z_j : \Omega \rightarrow \mathbb{C}$, $1 \leq j \leq m$, of class C^1 , satisfying:

- (1) If L is a local section of \mathcal{L} defined on Ω , then $LZ_j = 0$, $1 \leq j \leq m$;
- (2) $dZ_1(p) \wedge \cdots \wedge dZ_m(p) \neq 0$.

When \mathcal{L} and the first integrals Z_1, \dots, Z_m are of class C^∞ , the approximation theorem of Baouendi and Treves ([BT]) states that every distribution solution u of the sections of \mathcal{L} is locally the limit of a sequence of smooth solutions of the form $P_k \circ Z$ where $Z = (Z_1, \dots, Z_m)$ and the P_k are holomorphic polynomials. In this latter smooth category, the approximation theorem has had many consequences and has been a tool in dozens of papers, mainly dealing with CR theory (approximation of CR distributions, extension of CR distributions, propagation of analyticity), local behavior of solutions (local solvability, hypoellipticity, boundary behavior of solutions of homogeneous equations, uniqueness in the Cauchy problem) and miscellaneous topics like the similarity principle, the F. and M. Riesz theorem, representation of solutions, etc. To discuss uniqueness results, we recall that a submanifold $\Sigma \subseteq \mathcal{M}$ is called maximally real if for each $p \in \Sigma$,

$$\mathbb{C}T_p(\mathcal{M}) = \mathbb{C}T_p(\Sigma) \oplus \mathcal{L}_p.$$

In the CR case, that is, when $\mathcal{L} \cap \overline{\mathcal{L}} = 0$, Σ is maximally real if and only if it is totally real of maximal dimension. In the smooth case, if Σ is a maximally real submanifold and u is a distribution solution, the approximating functions $P_k \circ Z$ in the Baouendi-Treves approximation theorem are expressed in terms of the trace of u on Σ . Consequently, if u vanishes on Σ , then it vanishes in a neighborhood of Σ . In particular, if $\mathcal{N} \subseteq \mathcal{M}$ is a noncharacteristic hypersurface and a solution u vanishes

1991 *Mathematics Subject Classification*. Primary 35F15, 35B30, 42B30; Secondary 42A38, 30E25.

Work supported in part by NSF INT-0203005, CNPq and FAPESP.

on \mathcal{N} , then it vanishes in a neighborhood of \mathcal{N} since each $p \in \mathcal{N}$ is contained in a maximally real submanifold $\Sigma \subseteq \mathcal{N}$. The unique determination of a solution by its trace on a noncharacteristic hypersurface has the further consequence that the support of a solution propagates along the Sussmann orbits of $\text{Re } \mathcal{L} = \{X : X = \text{Re } L, L \text{ a smooth section of } \mathcal{L}\}$ ([T]). In this paper we explore the validity of the approximation theorem and the uniqueness results stated above for \mathcal{L} that is assumed to be only continuous with C^1 local first integrals. When \mathcal{L} is only continuous, we consider solutions u that are measures. In the smooth case, whenever Σ is a maximally real submanifold and u is a distribution solution, it is well known that u is a smooth function of variables transversal to Σ valued in the spaces of distributions on maximally real submanifolds. In particular, the trace of u on Σ —a key ingredient in the approximation theorem—is well defined. When \mathcal{L} is assumed to be only continuous and u is a measure solution, we show in section 2 that near each $p \in \mathcal{M}$, there is a nonvanishing continuous function a such that the trace of au on each noncharacteristic hypersurface \mathcal{N} is well defined. We also show that if the trace of au on \mathcal{N} vanishes, then u vanishes in a neighborhood of \mathcal{N} . This implies the propagation of the support of u along the orbits of $\text{Re } \mathcal{L}$. In section 3 we establish the approximation theorem for measure solutions. The results in this article extend some previous work done in [CR] (see Remark 2.6 for more on this).

2. REGULARITY AND UNIQUENESS FOR MEASURE SOLUTIONS

Since the problems we study are local, we will assume that \mathcal{M} is an open subset of \mathbb{R}^N . Consider a continuous vector subbundle \mathcal{L} of $\text{CT}(\mathcal{M})$ with fiber dimension $1 \leq n < N$ and set $m = N - n$.

Definition 2.1. *We say that \mathcal{L} is locally integrable if every $p \in \mathcal{M}$ is contained in an open set Ω such that there exist m functions $Z_j : \Omega \rightarrow \mathbb{C}$, $1 \leq j \leq m$, of class C^1 satisfying:*

- (1) *if L is a local section of \mathcal{L} defined on Ω , then $LZ_j = 0$, $1 \leq j \leq m$;*
- (2) *$dZ_1(p) \wedge \cdots \wedge dZ_m(p) \neq 0$.*

The functions Z_1, \dots, Z_m are called a complete set of first integrals. Notice that (1) and (2) imply that dZ_1, \dots, dZ_m span the orthogonal bundle $\mathcal{L}^\perp \subset \text{CT}^(\mathbb{R}^N)$.*

Let $p \in \mathcal{M}$. In a neighborhood Ω of p , we may find C^1 coordinates $x_1, \dots, x_m, t_1, \dots, t_n$ that vanish at p so that in these coordinates, a complete set of first integrals has the form

$$Z_j(x, t) = x_j + i\varphi_j(x, t), \quad j = 1, \dots, m,$$

with φ_j real satisfying $\partial\varphi_j/\partial x_k(0, 0) = 0$. By continuity, we may also assume that $|\partial\varphi_j/\partial x_k(0, 0)|$ remain small on Ω . In particular, we may assume that the square matrix $Z_x = [\partial Z_\ell/\partial x_k]$ is nonsingular and hence find continuous function $\lambda_{jk}(x, t)$ such that

$$\frac{\partial Z_\ell}{\partial t_j} + \sum_{k=1}^m \lambda_{jk} \frac{\partial Z_\ell}{\partial x_k} = 0, \quad j = 1, \dots, n, \quad \ell = 1, \dots, m,$$

which means that the vector fields

$$\frac{\partial}{\partial t_j} + \sum_{k=1}^m \lambda_{jk} \frac{\partial}{\partial x_k}, \quad j = 1, \dots, n,$$

are local generators of \mathcal{L} in Ω . It will be convenient to use another set of generators, namely,

$$L_j = \det Z_x \frac{\partial}{\partial t_j} + \det Z_x \sum_{k=1}^m \lambda_{jk} \frac{\partial}{\partial x_k}, \quad j = 1, \dots, n,$$

which have the advantage of being antisymmetric, that is,

$$\int (L_j u) v \, dx dt = - \int u L_j v \, dx dt, \quad j = 1, \dots, m, \quad u, v \in C_c^\infty(\Omega),$$

and may be written as partial differential operators in divergence form

$$L_j u = \frac{\partial(\det Z_x u)}{\partial t_j} + \sum_{k=1}^m \frac{\partial(\det Z_x \lambda_{jk} u)}{\partial x_k}, \quad u \in C_c^\infty(\Omega).$$

Thus, denoting by $\langle u, v \rangle = \int uv \, dx dt$ the usual duality pairing, if $u \in \mathcal{D}'_0(\Omega)$, we have that $L_j u$ may be defined by

$$\langle L_j u, v \rangle = - \langle u, L_j v \rangle, \quad v \in C_c^1(\Omega),$$

and $L_j u \in \mathcal{D}'_1(\Omega)$. The fact that u satisfies the set of equations

$$(2.2) \quad L_j u = 0, \quad j = 1, \dots, n,$$

has the following interpretation. Since we are interested in local questions, we may identify top forms $\psi(x) dx_1 \wedge \dots \wedge dx_N$ with functions $\psi(x)$, view a measure as acting continuously on top forms with continuous coefficients, and state

Definition 2.2. *We say the measure u is a solution of $\mathcal{L}u = 0$ if for each p and some complete set of C^1 first integrals Z_1, \dots, Z_m near p ,*

$$(2.3) \quad \langle u, dZ_1 \wedge \dots \wedge dZ_m \wedge d\omega \rangle = 0$$

for any ω which is an $(n-1)$ -form of class C^1 with compact support.

If we take

$$\omega = v dt_1 \wedge \dots \wedge \widehat{dt}_j \wedge \dots \wedge dt_n, \quad v \in C_c^1(\Omega),$$

where the hat means that the factor dt_j has been omitted, and express dv in the basis $dt_1, \dots, dt_n, dZ_1, \dots, dZ_m$, we see that

$$(2.4) \quad \begin{aligned} dZ_1 \wedge \dots \wedge dZ_m \wedge d\omega &= (-1)^{j-1} (\det Z_x)^{-1} L_j v dZ_1 \wedge \dots \wedge dZ_m \wedge dt_1 \wedge \dots \wedge dt_n \\ &= (-1)^{j-1} L_j v dx_1 \wedge \dots \wedge dx_m \wedge dt_1 \wedge \dots \wedge dt_n. \end{aligned}$$

Thus, using (2.4) for $j = 1, \dots, n$, it is easy to check that (2.2) and (2.3) are equivalent conditions for $u \in \mathcal{D}'_0(\Omega)$ in a neighborhood of p . Furthermore, we note that if (2.3) holds for some $\{Z_1, \dots, Z_m\}$, then it also holds for any other complete set of C^1 first integrals $\{W_1, \dots, W_m\}$. Indeed, by the approximation theorem (which is valid for C^1 functions) there exist holomorphic functions F_k^j such that for each $j = 1, \dots, n$, we may write $W_j = \lim_{k \rightarrow \infty} F_k^j(Z_1, \dots, Z_m)$ in the C^1 topology and so $dW_1 \wedge \dots \wedge dW_m = \lambda dZ_1 \wedge \dots \wedge dZ_m$ for some continuous function $\lambda = \lim_{k \rightarrow \infty} H_k(Z_1, \dots, Z_m)$ with H_k holomorphic. Since this process can be reversed, expressing $dZ_1 \wedge \dots \wedge dZ_m$ as a multiple of $dW_1 \wedge \dots \wedge dW_m$, it follows that $\lambda \neq 0$ on a neighborhood of p . Observe next that, since H_k is holomorphic, for any $(n-1)$ -form of class C^1 with compact support ω ,

$$H_k(Z) dZ_1 \wedge \dots \wedge dZ_m \wedge d\omega = dZ_1 \wedge \dots \wedge dZ_m \wedge d(H_k(Z)\omega)$$

and therefore,

$$\begin{aligned} \langle u, dW_1 \wedge \dots \wedge dW_m \wedge d\omega \rangle &= \lim_{k \rightarrow \infty} \langle u, H_k(Z) dZ_1 \wedge \dots \wedge dZ_m \wedge d\omega \rangle \\ &= \lim_{k \rightarrow \infty} \langle u, dZ_1 \wedge \dots \wedge dZ_m \wedge d(H_k(Z)\omega) \rangle = 0. \end{aligned}$$

Hence $\langle u, dW_1 \wedge \dots \wedge dW_m \wedge d\omega \rangle = 0$ for all $(n-1)$ -forms ω of class C^1 with compact support in a convenient neighborhood of p . Assume now that $n = 1$, so \mathcal{L} is locally spanned by a single vector field

$$L = \det Z_x \frac{\partial}{\partial t} + \det Z_x \sum_{k=1}^m \lambda_k \frac{\partial}{\partial x_k}$$

and let $u \in \mathcal{D}'_0(\Omega)$ satisfy the equation

$$(2.5) \quad Lu = 0$$

in the sense given by (2.3), so $\langle u, Lv \rangle = 0$, $v \in C_c^\infty(\Omega)$. We will prove that $Z_x(x, t)u$ may be identified with a function of t valued in $\mathcal{D}'(Q_R)$ for a convenient ball $Q_R \subset \mathbb{R}^m$. Since the statement is local it will be enough to restrict our attention to a neighborhood of the form $\Omega_R = Q_R \times I_R$, where I_R is the interval $(-R, R)$ and Q_R denotes a ball of radius R centered at the origin of \mathbb{R}^m . Given $\psi(x) \in C_c^\infty(Q_R)$, consider the distribution $u_\psi \in \mathcal{D}'(\mathbb{R})$ defined by

$$\langle u_\psi, \phi \rangle = \langle u, \psi \otimes \phi \rangle, \quad \phi \in C_c^\infty(I_R),$$

where $(\psi \otimes \phi)(x, t) = \psi(x)\phi(t)$. We first point out that $u_\psi \in \mathcal{D}'_0(I_R)$ for ψ fixed. Indeed, writing $U = u_\psi$,

$$|\langle U, \phi \rangle| = |\langle u, \psi \otimes \phi \rangle| \leq C \|\psi \otimes \phi\|_{C^0} \leq C'(R, \psi) \|\phi\|_{C^0}$$

showing that $U \in \mathcal{D}'_0(I_R)$. Set

$$V = (\det Z_x u)_\psi \quad \text{and} \quad W = - \sum_{k=1}^m (\det Z_x \lambda_k u)_{\psi x_j}.$$

Reasoning as before, we conclude that V and $W \in \mathcal{D}'_0(\mathbb{R})$; furthermore (2.5) implies that

$$\frac{dV}{dt} = W$$

holds in the sense of distributions. Thus, there exists a unique function $f_\psi(t)$, $t \in \mathbb{R}$, continuous from the right and, after shrinking I_R if necessary, of bounded variation on I_R , such that

$$\langle V, \phi \rangle = \int f_\psi(t) \phi(t) dt.$$

Since $\psi \mapsto f_\psi$ is linear and continuous in $C_c^1(I_R)$, we may write $f_\psi(t) = \langle f(t), \psi \rangle$ and consider $f(t)$ as an element of $\mathcal{D}'_1(Q_R)$ for each fixed t . If ψ belongs to a bounded subset \mathcal{B} of $C_c^1(\mathbb{R})$ and $|t| \leq R$, the variation of $f_\psi(t)$ is bounded by a constant depending only on \mathcal{B} and R . Thus

$$\langle \det Z_x u, \phi \otimes \psi \rangle = \int \langle f(t), \psi \rangle \phi(t) dt, \quad \phi(t) \in C_c^\infty(I_R), \psi(x) \in C_c^\infty(Q_R).$$

More generally, we have that

$$\langle \det Z_x u, \phi \rangle = \int \langle f(t), \phi(\cdot, t) \rangle dt, \quad \phi(x, t) \in C_c^\infty(\Omega_R).$$

Next for $\phi(x, t)$ a C^1 function we wish to understand the Leibnitz rule for

$$\frac{d}{dt} \langle f(t), \phi(\cdot, t) \rangle.$$

First fix $\psi(x) \in C_c^1(\mathbb{R}^m)$. Then we know that in the weak sense (which is what we will always mean)

$$\frac{d}{dt} \langle f(t), \psi(\cdot) \rangle = W_\psi(t)$$

where for $\varphi(t) \in C_c^0(\mathbb{R})$,

$$\langle W_\psi, \varphi(t) \rangle = \sum_k \iint (\lambda_k \det Z_x u) \frac{\partial \psi}{\partial x_k}(x) \varphi(t) dx dt.$$

We also know that there exists $C > 0$ such that $\forall \psi(x) \in C_c^1(B_r)$, and $|t| \leq R$,

$$(2.6) \quad |\langle f(t), \psi \rangle| \leq C \|\psi\|_{C^1}.$$

Let $\phi(x, t) \in C_c^1$. Consider $F(t) = \langle f(t), \phi(\cdot, t) \rangle$. We will show that $F(t)$ is a function of bounded variation and write a formula for its weak derivative. Clearly, $F(t)$ is a measurable function. From (2.6) we see that $F(t)$ is a bounded function and so it defines a distribution. Suppose now $\phi(x, t) = \psi(x)\phi(t) \in C_c^1$. Then

$$\frac{d}{dt} \langle f(t), \phi(\cdot, t) \rangle = \phi'(t) \langle f(t), \psi(\cdot) \rangle + \phi(t) W_\psi$$

and so for any $h(t) \in C_c^0$,

$$(2.7) \quad \left(\frac{d}{dt} \langle f(t), \phi(\cdot, t) \rangle \right) (h) = \iint \det Z_x u \psi(x) \phi'(t) h(t) dx dt \\ + \sum_{k=1}^m \iint (\lambda_k \det Z_x u) \frac{\partial \psi}{\partial x_k}(x) \phi(t) h(t) dx dt.$$

More generally, we claim that for any $\phi(x, t) \in C_c^1$, the formula

$$(2.8) \quad \frac{d}{dt} \langle f(t), \phi(\cdot, t) \rangle = (\det Z_x u)_{\phi_t} + \sum_{k=1}^m (\lambda_k \det Z_x u)_{\phi_{x_k}}$$

holds where in the right hand side, $(\det Z_x u)_{\phi_t}$ is the distribution in t space defined by

$$\langle (\det Z_x u)_{\phi_t}, h(t) \rangle = \iint (\det Z_x u)(x, t) \phi_t(x, t) h(t) dx dt$$

and likewise for $\sum_k (\lambda_k \det Z_x u)_{\phi_{x_k}}$. Equation (2.7) shows that (2.8) easily holds for C^1 functions of the form

$$\sum_{j=1}^N \sum_{k=1}^M \psi_j(x) \phi_k(t).$$

The general case follows by approximating a given C^1 function $\phi(x, t)$ by such finite sums and observing that each side of (2.8) is a continuous operator on the space of C^1 functions. Observe that (2.8) in particular shows that when $\phi(x, t)$ is a C^1 function, $\frac{d}{dt} \langle f(t), \phi(\cdot, t) \rangle$ is a measure and hence $\langle f(t), \phi(\cdot, t) \rangle$ is a function of bounded variation.

Recall that for every $t_0 \in \mathbb{R}$, $|t_0| < R$, $\det Z_x u$ has a trace at $t = t_0$ which is a distribution of order one. This trace is given by $f(t_0)$ and satisfies the one-sided continuity property $\lim_{\varepsilon \searrow 0} f(t_0 + \varepsilon) = f(t_0)$. We may also denote this trace by $Z_x(\cdot, t_0)u(\cdot, t_0)$ or $Z_x(x, t_0)u(x, t_0)$ rather than $f(t_0)$. We now recall the basic operators in the Baouendi-Treves approximation formula, namely

$$E_\tau u(x, t) = (\tau/\pi)^{m/2} \int_{\mathbb{R}^m} e^{-\tau[Z(x,t)-Z(x',0)]^2} \det Z_x(x', 0) u(x', 0) h(x') dx', \\ G_\tau u(x, t) = (\tau/\pi)^{m/2} \int_{\mathbb{R}^m} e^{-\tau[Z(x,t)-Z(x',t)]^2} \det Z_x(x', t) u(x', t) h(x') dx', \\ R_\tau u(x, t) = G_\tau u(x, t) - E_\tau u(x, t),$$

where the notation $[Z(x, t) - Z(x', t')]^2$ means $\sum_{j=1}^m (Z_j(x, t) - Z_j(x', t'))^2$ and the function $h(x) \in C_c^\infty(Q_R)$ is supported in a conveniently small interval and it is identically equal to 1 on a neighborhood of the origin. In the classical case, when L and $Z(x, t)$ are smooth and u is a distribution solution of $Lu = 0$, it can be proved that $R_\tau u(x, t) \rightarrow 0$ as $\tau \rightarrow \infty$ on a small neighborhood of the origin, with convergence in the C^∞ topology. We now adapt some of the classical arguments to

the present situation, where $u \in \mathcal{D}'_0$ and \mathcal{L} is of class C^0 , to show that $R_\tau u(x, t) \rightarrow 0$ uniformly as $\tau \rightarrow \infty$ on some neighborhood of the origin. Write, for $\zeta \in \mathbb{C}^m$ and $t \in I_{\mathbb{R}}$,

$$\begin{aligned} \tilde{G}_\tau u(\zeta, t) &= \left\langle f(t), e^{-\tau[\zeta - Z(\cdot, t)]^2} h(\cdot) \right\rangle = \left\langle \det Z_x(\cdot, t) u(\cdot, t), e^{-\tau[\zeta - Z(\cdot, t)]^2} h(\cdot) \right\rangle \\ &= \int_{\mathbb{R}^m} e^{-\tau[\zeta - Z(x', t)]^2} \det Z_x(x', t) u(x', t) h(x') dx', \end{aligned}$$

and

$$\tilde{R}_\tau u(\zeta, t) = \tilde{G}_\tau u(\zeta, t) - \tilde{G}_\tau u(\zeta, 0),$$

so we have

$$R_\tau u(x, t) = (\tau/\pi)^{m/2} \tilde{R}_\tau u(Z(x, t), t).$$

If $F : I_{\mathbb{R}} \rightarrow \mathbb{R}$ is a function of bounded variation with weak derivative $F' = \mu$, the fundamental theorem of calculus for F states that

$$\mu((a, b]) = \int_{(a, b]} d\mu = F(b+) - F(a+), \quad -R < a < b < R.$$

Let $F(t) = \tilde{G}_\tau u(\zeta, t)$. Since the function $e^{-\tau[\zeta - Z(x, t)]^2} h(x)$ is C^1 , we have seen that $F(t)$ is a function of bounded variation, continuous from the right, and formula (2.8) leads to

$$\begin{aligned} \tilde{R}_\tau u(\zeta, t) &= F(t) - F(0) \\ &= \sum_{k=1}^m \int_0^t \int_{\mathbb{R}^m} \lambda_k(x', t') \det Z_x u(x', t') \partial_{x'_k} (h(x') e^{-\tau[\zeta - Z(x', t')]^2}) dx' dt' \\ &\quad + \int_0^t \int_{\mathbb{R}^m} (\det Z_{x'} u)(x', t') h(x') \partial_{t'} (e^{-\tau[\zeta - Z(x', t')]^2}) dx' dt' \end{aligned}$$

where \int_0^t must be understood as $\int_{(0, t]}$ for $t \geq 0$ and as $-\int_{(t, 0]}$ for $t < 0$. Using the fact that

$$\left(\partial_{t'} + \sum_{k=1}^m \lambda_k \partial_{x'_k} \right) e^{-\tau[\zeta - Z(x', t')]^2} = 0$$

we get

$$(2.6) \quad \tilde{R}_\tau u(\zeta, t) = \int_0^t \int_{\mathbb{R}^m} u(x', t') e^{-\tau[\zeta - Z(x', t')]^2} Lh(x', t') dx' dt'$$

which, for $\zeta = Z(x, t)$ gives

$$R_\tau u(x, t) = (\tau/\pi)^{m/2} \int_0^t dt' \int_{\mathbb{R}^m} e^{-\tau[Z(x, t) - Z(x', t')]^2} u(x', t') Lh(x', t') dx',$$

which extends the classical expression for $R_\tau u$. If $\delta > 0$ is chosen so that $h(x') \equiv 1$ for $|x'| \leq 2\delta$, and we take $|x| \leq \delta$ and t sufficiently small, the exponential in the

expression of $R_\tau u(x, t)$ can be majorized by $e^{-c\tau}$ for some $c > 0$, allowing us to conclude that the estimate $|R_\tau u(x, t)| \leq Ce^{-c\tau}$ holds in a neighborhood Ω of the origin and, in particular, $R_\tau u(x, t) \rightarrow 0$ uniformly on Ω as $\tau \rightarrow \infty$. On the other hand, standard arguments show that, $G_\tau u(x, t)$ converges weakly to the measure u . It follows that $E_\tau u(x, t) \rightarrow u(x, t)$ in $\mathcal{D}'_0(\Omega)$ as $\tau \rightarrow \infty$. If we assume now that the trace $\det Z_x(x, 0)u(x, 0) \equiv 0$, we see that $E_\tau u(x, t) \equiv 0$ showing that $u = 0$ on Ω . Suppose next the left hand limit $\lim_{t \rightarrow 0^-} Z_x(x, t)u(x, t) = (Z_x u)(x, 0^-) = 0$. Then the equation $\mu([b, 0)) = F(0^-) - F(b^-)$ for $b < 0$ implies that $(Z_x u)(x, b^-) = 0$ for $b < 0$. Since F is continuous except possibly on a countable set, we can get $b < 0$ arbitrarily close to 0 where F is continuous and hence where the trace $Z_x(x, b)u(x, b) = 0$. We can then use the approximation scheme as above by using the trace at such b (with b sufficiently small) in the definition of the operators $E_\tau u(x, t)$ to conclude that $u(x, t) \equiv 0$ in a neighborhood of the origin. This shows that a function of bounded variation $f(t)$ satisfying

$$\langle \det Z_x u, \phi \otimes \psi \rangle = \int \langle f(t), \psi \rangle \phi(t) dt, \quad \phi(t) \in C_c^\infty(I_R), \psi(x) \in C_c^\infty(Q_R),$$

cannot have a jump at point $t = t_0$ unless both $f(t_0+) \neq 0$ and $f(t_0-) \neq 0$. Notice that the value $f(t)$ is completely determined outside the at most countable set at which jumps occur and that at the jumps the value of f is arbitrary, unless one imposes, say, that f is continuous from the right or continuous from the left. We may now state and prove

Theorem 2.3. *Let \mathcal{L} be a locally integrable continuous subbundle of $\mathcal{CT}(\mathcal{M})$. Let $u \in \mathcal{D}'_0$ be a Radon measure such that $\mathcal{L}u = 0$, and let Σ be a C^1 hypersurface noncharacteristic with respect to \mathcal{L} . Let $p \in \Sigma$ and let $N = N(p)$ be a unit vector normal to Σ at p . There exists a nonvanishing continuous function D defined in a neighborhood Ω of p such that $u_1 = Du$ has a well defined trace $T_\tau u_1$ defined on the translations $(\tau N + \Sigma) \cap \Omega$ along N satisfying the following properties:*

- (1) $\lim_{\varepsilon \searrow 0} T_{\tau+\varepsilon} u_1 = T_\tau u_1$ (one-sided continuity);
- (2) if $T_0 u_1 = 0$ on $\Sigma \cap \Omega$ then u vanishes identically on a neighborhood of p (uniqueness).

Proof. We may find local C^1 coordinates $x_1, \dots, x_m, t_1, \dots, t_n$, defined in a neighborhood Ω of p so that $x_k(p) = t_j(p) = 0$, $1 \leq j \leq n$, $1 \leq k \leq m$, and a complete set of first integrals is of the form

$$Z_j(x, t) = x_j + i\varphi_j(x, t), \quad j = 1, \dots, m,$$

with $\partial\varphi_j/\partial x_k(0, 0) = 0$. Furthermore, we may assume that $\Sigma \cap \Omega$ is given by $t_1 = 0$ and that N points in the direction of increasing t_1 . If $n = 1$, we have just seen that taking $D = \det Z_x$ the conclusions of the theorem are obtained. For $n > 1$ we discard the vector fields L_2, \dots, L_n and rename the coordinates as follows: $t' = t_1$, $x'_k = x_k$, $1 \leq k \leq m$, $x'_{m+k} = t_{k+1}$, $1 \leq k \leq n-1$. Then we add new first integrals $Z_{m+1}(x', t') = x'_{m+1}, \dots, Z_{m+n-1}(x', t') = x'_{m+n-1}$. This leads us to a new locally integrable structure $\tilde{\mathcal{L}}$ of dimension 1 spanned by $L = L_1$ such that

$\tilde{\mathcal{L}}u = 0$. Hence, we may reason as before and conclude that $\det Z_x u$ has a trace on Σ with the required properties. \square

Note that we may as well define a trace that possesses the uniqueness property (2) but instead of (1) satisfies $\lim_{\varepsilon \searrow 0} T_{\tau-\varepsilon} u_1 = T_\tau u_1$.

If u vanishes as a distribution above a noncharacteristic hypersurface Σ it is clear that $T_\tau u_1 = 0$ for $\tau > 0$ and sufficiently small. Then the continuity property (1) of the trace $T_\tau u_1$ implies that $T_0 u_1 = 0$, so the support of u cannot meet Σ by property (2). If u vanishes below Σ we reach the same conclusion by using the trace that is continuous from the left. It follows from Theorem 2.3 that the support of u propagates along the normal of noncharacteristic hypersurfaces, which implies in a standard way that the support of u is a union of orbits of \mathcal{L} .

Corollary 2.4. *Let \mathcal{L} be a locally integrable continuous subbundle of $\mathbb{C}T(\Omega)$, $\Omega \subset \mathbb{R}^N$ open. Let $u \in \mathcal{D}'_0(\Omega)$ be a Radon measure such that $\mathcal{L}u = 0$. Then the support of u is a union of orbits of \mathcal{L} in Ω .*

For the case of a CR structure on a generic CR submanifold $M \subset \mathbb{C}^n$ of class C^1 , we recover a result of [CR]:

Corollary 2.5. ([CR]) *Let M be a generic CR submanifold of \mathbb{C}^n of class C^1 . Then the support of a CR distribution $u \in \mathcal{D}'_0(M)$ is a union of orbits.*

Remark 2.6. In the CR case, there are two differences between our results here and the ones proved in the paper [CR]. First, in [CR], Corollary 2.5 is deduced from a weaker uniqueness result (Lemma A.5 in [CR]) than the one stated in part 2 of Theorem 2.3. Second, the results in [CR] do not show that a measure solution can be approximated by C^1 solutions as is done in section 3 here (see Theorem 3.2).

Remark 2.7. If μ is a Radon measure that has a trace defined at a family of translations of a hypersurface Σ and f is a continuous function, it is not true, in general, that the product $f\mu$ also has a trace. For instance, the function $\mu(x, t) = 1/(x + it) \in L^1_{\text{loc}}(\mathbb{R}^2)$ has a trace $T_t \mu$, continuous from the right, given by the locally integrable function $\mathbb{R} \ni x \mapsto 1/(x + it)$ for $t \neq 0$ and equal to the distribution $T_0 \mu = \text{pv}1/x - i\pi\delta$ for $t = 0$. Consider the continuous function defined by $f(x, t) = 0$ for $x \leq 0$ and $f(x, t) = 1/|\ln x|$ for $x > 0$ and set $\nu(x, t) = (f\mu)(x, t)$. Thus, $\nu \in L^1_{\text{loc}}(\mathbb{R}^2)$ is continuous off the origin and has a natural restriction to the straight lines $t = \text{constant} \neq 0$ given by restriction. On the other hand, if $v(x) \in C_c^\infty(\mathbb{R})$ is real and $v(0) \neq 0$ we see that

$$\lim_{t \rightarrow 0} \text{Re} \int \nu(x, t)v(x) dx = \int_0^\infty \frac{v(x)}{x|\ln x|} dx = \infty,$$

so it is not possible to define a one-sided continuous trace for ν that coincides with restriction for $t \neq 0$.

3. APPROXIMATION OF MEASURE SOLUTIONS BY SMOOTHER SOLUTIONS.

Assume as in section 2 that \mathcal{L} is a continuous locally integrable structure defined in an open subset Ω of \mathbb{R}^N that contains the origin, with fiber dimension n , and

that $u \in \mathcal{D}'_0(\Omega)$ satisfies the equation $\mathcal{L}u = 0$. Shrinking Ω we may assume that u is a finite measure and choosing appropriate C^1 coordinates $x_1, \dots, x_m, t_1, \dots, t_n$, we may assume that a complete set of first integrals are of the form

$$Z_j(x, t) = x_j + i\varphi_j(x, t), \quad j = 1, \dots, m,$$

with φ_j real satisfying $\partial\varphi_j/\partial x_k(0, 0) = 0$ and $|\partial\varphi_j/\partial x_k(0, 0)|$ small throughout Ω . Consider balls $B_1 \subset \mathbb{R}^m$, $B_2 \subset \mathbb{R}^n$, $B_1 = B(0, r_1)$, $B_2 = B(0, r_2)$, such that $B_1 \times B_2 \subset \Omega$, and for $\psi(x) \in C_c^1(B_1)$ set

$$\langle (\det Z_x u)_\psi, \phi \rangle = \langle \det Z_x u, \psi \otimes \phi \rangle, \quad \phi(t) \in C_c^1(B_2).$$

It follows from the discussion in section 2 that $(\det Z_x u)_\psi$ is a finite measure on B_2 such that its exterior derivative $d_t(\det Z_x u)_\psi$ is also a finite measure. In other words, $(\det Z_x u)_\psi$ may be identified with a function of bounded variation $f_\psi(t) = \langle f(t), \psi \rangle \in BV(W)$, $\bar{B}_2 \subset W$. We now recall some general properties of functions of bounded variation (see, e.g., [G]): if $g(t) \in BV(W)$ then $g(t) \in L^1(W)$ and its derivatives $\partial g/\partial t_j = \mu_j$ are measures with total bounded variation $\|\mu_j\| = |\mu_j|(W)$, where $|\mu|$ is the variation of μ . In fact, $g(t)$ belongs to the smaller space $L^p(B_2)$, with $p = p(n) = n/(n-1)$, and the generalized Poincaré inequality holds: if we denote by

$$g_{B_2} = \frac{1}{|B_2|} \int_{B_2} g(t) dt$$

the mean value of g on B_2 , then

$$\|g - g_{B_2}\|_{L^p(B_2)} \leq C \sum_{j=1}^n |\mu_j|(B_2),$$

where $C > 0$ depends on the dimension n but neither on the radius of B_2 nor on g . A pointwise estimate related to the Poincaré inequality is given in the following lemma.

Lemma 3.1. *Let g be of bounded variation on an open set $W \subset \mathbb{R}^n$ that contains \bar{B}_2 and set $\partial g/\partial t_j = \mu_j$, $|\mu| = \sum_{j=1}^n |\mu_j|$. Then*

$$(3.1) \quad |g(t) - g_{B_2}| \leq C \int_{B_2} |t - s|^{1-n} d|\mu|(s), \quad a.e. \ t \in B_2$$

where C depends only on n .

Proof. For $g \in W^{1,1}(B_2)$ the lemma is well known (see, e.g., [GT, Lemma 7.16]). Let g be as in the lemma. Let $\eta(s) \in C_c^\infty(\mathbb{R}^n)$ such that $\eta(s)$ is nonnegative, its support lies in the unit ball $B_1(0)$ and $\int_{\mathbb{R}^n} \eta(s) ds = 1$. Set $\eta_\epsilon(s) = \epsilon^{-n} \eta(s/\epsilon)$. If

$$g_\epsilon(t) = \eta_\epsilon * g(t),$$

then $g_\epsilon \rightarrow g$ in L^1 and $g_\epsilon(t) \rightarrow g(t)$ a.e. For $t \in B_2$, and for any $\epsilon > 0$, we have

$$(3.2) \quad |g_\epsilon(t) - (g_\epsilon)_{B_2}| \leq C \int_{B_2} \frac{|Dg_\epsilon(s)|}{|s - t|^{n-1}} ds.$$

Note that

$$Dg_\epsilon(s) = \eta_\epsilon * \mu(s)$$

where $\mu = (\mu_1, \dots, \mu_n)$. It follows from (3.2) that

$$(3.3) \quad |g_\epsilon(t) - (g_\epsilon)_{B_2}| \leq C \int_{B_{r_2+\epsilon}} \left(\int_{B_2} \frac{\eta_\epsilon(s' - s)}{|s - t|^{n-1}} ds \right) d|\mu|(s').$$

We will estimate the inner integral which after a change of variables is dominated by

$$\int_{|s| \leq 1} \frac{1}{|s' - t - \epsilon s|^{n-1}} ds.$$

For $y \in \mathbb{R}^n$ fixed, $y \neq 0$, consider therefore the integral

$$(3.4) \quad \int_{|s| \leq 1} \frac{1}{|y - \epsilon s|^{n-1}} ds.$$

If $|y| \geq 2\epsilon$, then for all $|s| \leq 1$, $|y - \epsilon s| \geq \frac{|y|}{2}$ and so for such y ,

$$\int_{|s| \leq 1} \frac{1}{|y - \epsilon s|^{n-1}} ds \leq \frac{c_n}{|y|^{n-1}}.$$

Suppose $|y| \leq 2\epsilon$. In the integral in (3.4), change variables $t' = \epsilon s$. We then get

$$\begin{aligned} \int_{|s| \leq 1} \frac{1}{|y - \epsilon s|^{n-1}} ds &= \frac{1}{\epsilon^n} \int_{|t'| \leq \epsilon} \frac{1}{|y - t'|^{n-1}} dt' \\ &\leq \frac{1}{\epsilon^n} \int_{|t'| \leq 3\epsilon} \frac{1}{|t'|^{n-1}} dt' \\ &\leq \frac{c}{\epsilon^{n-1}} \leq \frac{c_n}{|y|^{n-1}}. \end{aligned}$$

It now follows that for some dimensional constant C_n ,

$$(3.5) \quad \int_{B_2} \frac{\eta_\epsilon(s' - s)}{|s - t|^{n-1}} ds \leq \int_{|s| \leq 1} \frac{1}{|s' - t - \epsilon s|^{n-1}} ds \leq \frac{C_n}{|s' - t|^{n-1}}.$$

To prove (3.1) for a fixed $t \in B_2$, we may assume —and we will do so— that the function $s \mapsto |t - s|^{1-n} \in L^1(B_2, d|\mu|)$, otherwise the right hand side would be infinity and the inequality trivial. By the Dominated Convergence Theorem, it follows that

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \int_{B_{r_2+\epsilon}} \left(\int_{B_2} \frac{\eta_\epsilon(s' - s)}{|s - t|^{n-1}} ds \right) d|\mu|(s') \\ &= \int_{s'} \lim_{\epsilon \rightarrow 0} \chi_{B_{r_2+\epsilon}}(s') \left(\int_{B_2} \frac{\eta_\epsilon(s' - s)}{|s - t|^{n-1}} ds \right) d|\mu|(s') \\ &= \int_{\bar{B}_2} \left(\int_{t'} \frac{\eta(t')}{|s' - t|^{n-1}} dt' \right) d|\mu|(s') \\ &= \int_{\bar{B}_2} \frac{1}{|s' - t|^{n-1}} d|\mu|(s') \end{aligned}$$

where $\chi_{B_{r_2+\epsilon}}(s')$ denotes the characteristic function of the ball $B_{r_2+\epsilon}$. Thus, letting $\epsilon \searrow 0$ in (3.3), we obtain estimate (3.1) with the integral on the right hand side taken over \bar{B}_2 instead of B_2 . But then we obtain (3.1) by expressing $B_2 = B(0, r_2)$ as an increasing countable union of closed balls of radius $< r_2$ and applying the weaker estimate to the smaller closed balls. This proves the lemma. \square

Remark. It is known that (3.2) holds with $C = \gamma_n = 2^{n-1}\Gamma(n/2)/\pi^{n/2}$. Then the trivial estimate $|Dg_\epsilon(s)| \leq \sum_{j=1}^n |D_j g_\epsilon(s)|$ implies that we may take $C = \gamma_n$ in (3.3) and the proof of the lemma shows that (3.1) is valid with $C = \gamma_n$.

We will now introduce a variation of the Baouendi-Treves approximation formula that does not require a continuous trace of the solution to maximally real submanifolds. For $(x, t) \in B_1 \times B_2$, $\tau \geq 1$, and a fixed bump function $h(x) \in C_c^\infty(B_1)$ that is identically 1 on a neighborhood of the origin, set

$$F_\tau u(x, t) = (\tau/\pi)^{m/2} \frac{1}{|B_2|} \int_{\mathbb{R}^m \times B_2} e^{-\tau[Z(x,t)-Z(y,s)]^2} \det Z_y(y, s) u(y, s) h(y) dy ds.$$

The integral

$$\int_{\mathbb{R}^m \times B_2} e^{-\tau[Z(x,t)-Z(y,s)]^2} \det Z_y(y, s) u(y, s) h(y) dy ds$$

in the definition of $F_\tau u(x, t)$ is understood as

$$\int_{B_2} \left\langle f(s), e^{-\tau[Z(x,t)-Z(\cdot, s)]^2} h(\cdot) \right\rangle ds.$$

It is clear that, for fixed $\tau \geq 1$, $F_\tau u(x, t)$ is an entire function of $Z(x, t)$, in particular, it satisfies the equation $\mathcal{L}E_\tau u = 0$ in the classical sense and is of class C^1 . We wish to show that, for r_2 sufficiently small, $F_\tau u(x, t) \rightarrow u(x, t)$ in the weak-* topology of $C_c^0(U)^*$ as $\tau \rightarrow \infty$ for a convenient neighborhood of the origin $U \subset \mathbb{R}^m \times \mathbb{R}^n$. More precisely, we will show that, given $v \in C_c^0(U)$,

$$\langle u - F_\tau u, v \rangle \longrightarrow 0 \quad \text{as } \tau \rightarrow \infty.$$

For $s \in B_2$ and $\zeta \in \mathbb{C}^m$ consider the function

$$g^\tau(s, \zeta) = (\tau/\pi)^{m/2} \int_{\mathbb{R}^m} e^{-\tau[\zeta - Z(y, s)]^2} \det Z_y(y, s) u(y, s) h(y) dy.$$

Notice that $g^\tau(s, \zeta)$ is of the form $(\det Z_x u)_\psi$ for a convenient ψ depending on τ and ζ so $g^\tau(s, \zeta)$ is defined for all $\tau \geq 1$, all $\zeta \in \mathbb{C}$ and all $s \in B_2 \setminus E$, with $|E| = 0$, and belongs to $L^p(B_2)$ for τ, ζ fixed. To see that E can be taken independently of ζ and τ , we may replace the exponential by its Taylor series, obtaining a power series in ζ and τ with coefficients that are linear combinations of functions of the form

$$c_\alpha(s) = \int_{\mathbb{R}^m} Z(y, s)^\alpha \det Z_y(y, s) u(y, s) h(y) dy, \quad \alpha \in \mathbb{Z}_+^n,$$

defined for $s \in B_2 \setminus E_\alpha$, $|E_\alpha|=0$. Then we take $E = \bigcup_\alpha E_\alpha$. The norm of $c_\alpha(s)$ in $BV(B_2)$ is dominated by the C^1 norm of $Z(y, s)^\alpha h(y)$, which has polynomial growth in α , so the argument shows that we may regard $(\zeta, \tau) \mapsto g^\tau(s, \zeta)$ as an entire holomorphic function with values in $BV(B_2)$. Using (3.1) we get

$$(3.6) \quad |g^\tau(t, \zeta) - g_{B_2}^\tau(\zeta)| \leq C \int_{B_2} |t - s|^{1-n} d|\mu^\tau|(s, \zeta), \quad \text{a.e. } t \in B_2.$$

For $\zeta = Z(x, t)$ we get

$$\begin{aligned} g^\tau(t, Z(x, t)) &= G_\tau u(x, t) \\ g_{B_2}^\tau(Z(x, t)) &= F_\tau u(x, t) \end{aligned}$$

so (3.6) yields

$$(3.7) \quad |G_\tau u(x, t) - F_\tau u(x, t)| \leq C \int_{B_2} |t - s|^{1-n} d|\mu^\tau|(s, Z(x, t)),$$

for a.e. $t \in B_2$. To take advantage of estimate (3.7) we need to compute

$$\begin{aligned} \mu_j^\tau(t, \zeta) &= \frac{\partial}{\partial t_j} g^\tau(t, \zeta) \\ &= \frac{\partial}{\partial t_j} \int_{\mathbb{R}^m} e^{-\tau[\zeta - Z(y, t)]^2} \det Z_x(y, t) u(y, t) h(y) dy. \end{aligned}$$

Differentiating under the integral sign, using the equation $L_j u = 0$ in order to replace $\partial_{t_j}(\det Z_x u)$ by $-\sum_{k=1}^m \partial_{y_k}(\det Z_x \lambda_{jk} u)$ in the first term on the right, integrating by parts with respect to y and using the fact that

$$\left(\partial_{t_j} + \sum_{k=1}^m \lambda_{jk} \partial_{y_k} \right) e^{-\tau[\zeta - Z(y, t)]^2} = 0$$

we get

$$\begin{aligned} \mu_j^\tau(t, \zeta) &= \int_{\mathbb{R}^m} e^{-\tau[\zeta - Z(y, t)]^2} (\det Z_x u)(y, t) \sum_{k=1}^m \lambda_{jk} \frac{\partial h(y)}{\partial y_k} dy. \\ &= \int_{\mathbb{R}^m} e^{-\tau[\zeta - Z(y, t)]^2} (\det Z_x u)(y, t) h_j(y, t) dy. \end{aligned}$$

To estimate the variation $|\mu_j^\tau(\zeta)|$ of $\mu_j^\tau(\cdot, \zeta)$ on B_2 observe that if $\phi(t) \in L^\infty(B_2)$,

$$\int \mu_j^\tau(t, \zeta) \phi(t) dt = \int e^{-\tau[\zeta - Z(y, t)]^2} \det Z_x(y, t) h_j(y, t) \phi(t) du(y, t)$$

so, for any Borel set $E \subset B_2$,

$$\int_E d|\mu_j^\tau|(\zeta) \leq C \sup_{(y, s) \in \text{supp } dh \times B_2} e^{-\tau \Re[\zeta - Z(y, s)]^2} \int_{B_1 \times E} d|u|$$

where $|u|$ denotes the variation of u . For $\zeta = Z(x, t)$ we get

$$(3.8) \quad \int_E d|\mu_j^\tau|(Z(x, t)) \leq C \sup_{\{2\delta \leq |y| < r_1\} \times B_2} e^{-\tau \Re[Z(x, t) - Z(y, s)]^2} \int_{B_1 \times E} d|u|$$

if we assume that $h(y)$ has been chosen to satisfy $h(y) \equiv 1$ for $|y| \leq 2\delta < r_1$. Let $v(x, t) \in C_c^0(B(0, \delta) \times B_2)$ and consider

$$I_\tau(v) = \int (G_\tau u(x, t) - F_\tau u(x, t)) v(x, t) dx dt.$$

If (x, t) is in the support of $v(x, t)$ and r_2 is chosen conveniently small (depending on δ) we verify that, in the exponent on the right hand side of (3.8), $\Re[Z(x, t) - Z(y, s)]^2 \geq c > 0$. Thus, in view of (3.7) and (3.8),

$$\begin{aligned} |I_\tau(v)| &\leq C \|v\|_{L^\infty} e^{-c\tau} \int_{B_1 \times B_2} \int_{B_1 \times B_2} \frac{dx dt}{|t - s|^{n-1}} d|u|(y, s) \\ &\leq C \|v\|_{L^\infty} |u|(B_1 \times B_2) e^{-c\tau}. \end{aligned}$$

This shows that $I_\tau(v) \rightarrow 0$ as $\tau \rightarrow \infty$. On the other hand, adapting the arguments in [HM] (where $Z(x, t)$ is smooth) there is no difficulty in proving that, since $u(x, t)$ is a bounded measure, there is a neighborhood U of the origin in $\mathbb{R}^m \times \mathbb{R}^n$ such that $G_\tau u \rightarrow u$ in the weak-* topology of $C_c^0(U)^*$. Thus,

$$\langle u - F_\tau u, v \rangle \longrightarrow 0 \quad \text{as } \tau \rightarrow \infty,$$

as we wished to prove.

Remark. Integrating (3.7) with respect to $dx dt$ on $B_1 \times B_2$ and taking account of (3.8) we see that

$$\|G_\tau u - F_\tau u\|_{L^1(U)} \leq C, \quad \tau \geq 1.$$

Since, it is known that $\|G_\tau u\|_{L^1(U)} \leq C$ it follows that $\|F_\tau u\|_{L^1(U)} \leq C$ as well.

We have proved:

Theorem 3.2. *Let \mathcal{L} be a locally integrable continuous subbundle of $\mathcal{CT}(\mathcal{M})$. Let $u \in \mathcal{D}'_0$ be a Radon measure such that $\mathcal{L}u = 0$. Then for each $p \in \mathcal{M}$, there is a neighborhood Ω and a sequence P_k of holomorphic polynomials such that $u = \lim_{k \rightarrow \infty} P_k(Z)$ in the weak-* topology of $C_c^0(\Omega)^*$ where $Z = (Z_1, \dots, Z_m)$ is a complete set of first integrals on Ω .*

REFERENCES

- [BT] M. S. Baouendi and F. Trèves, *A property of the functions and distributions annihilated by a locally integrable system of complex vector fields*, Ann. of Math. **113** (1981), 387–421.
- [CR] E. M. Chirka and C. Rea, *Differentiable CR mappings and CR orbits*, Duke Math. J. **94** (1998), 325–340.
- [GT] D. Gilbard and N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag Berlin Heidelberg New York, 2001.
- [G] E. Giusti, *Minimal surfaces and functions of bounded variation*, Birkhäuser Boston, 1984.

- [HM] J. Hounie and P. Malagutti, *On the convergence of the Baouendi-Treves approximation formula*, Comm. P.D.E. **23** (1998), 1305–1347.
- [T] F. Treves, *Hypo-analytic structures, local theory*, Princeton University Press, 1992.

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA, PA 19122-6094,
USA

E-mail address: berhanu@temple.edu

DEPARTAMENTO DE MATEMÁTICA, UFSCAR, 13.565-905, SÃO CARLOS, SP, BRASIL

E-mail address: hounie@dm.ufscar.br